


## SECTION 1


- This section contains **SIX (06)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:
  - Full Marks* : +4 If only (all) the correct option(s) is(are) chosen;
  - Partial Marks* : +3 If all the four options are correct but **ONLY** three options are chosen;
  - Partial Marks* : +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
  - Partial Marks* : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
  - Zero Marks* : 0 If unanswered;
  - Negative Marks* : -2 In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to correct answers, then
  - choosing **ONLY** (A), (B) and (D) will get +4 marks;
  - choosing **ONLY** (A) and (B) will get +2 marks;
  - choosing **ONLY** (A) and (D) will get +2 marks;
  - choosing **ONLY** (B) and (D) will get +2 marks;
  - choosing **ONLY** (A) will get +1 mark;
  - choosing **ONLY** (B) will get +1 mark;
  - choosing **ONLY** (D) will get +1 mark;
  - choosing no option(s) (i.e. the question is unanswered) will get 0 marks and
  - choosing any other option(s) will get -2 marks.

Q.1 The reaction sequence(s) that would lead to *o*-xylene as the major product is(are)


(A)




(B)




(C)



(D)




Q.2 Correct option(s) for the following sequence of reactions is(are)



- (A) **Q** = KNO<sub>2</sub>, **W** = LiAlH<sub>4</sub>                      (B) **R** = benzenamine, **V** = KCN  
 (C) **Q** = AgNO<sub>2</sub>, **R** = phenylmethanamine    (D) **W** = LiAlH<sub>4</sub>, **V** = AgCN

Q.3 For the following reaction




the rate of reaction is  $\frac{d[\mathbf{P}]}{dt} = k[\mathbf{X}]$ . Two moles of **X** are mixed with one mole of **Y** to make 1.0 L of solution. At 50 s, 0.5 mole of **Y** is left in the reaction mixture. The correct statement(s) about the reaction is(are)

(Use:  $\ln 2 = 0.693$ )

- (A) The rate constant,  $k$ , of the reaction is  $13.86 \times 10^{-4} \text{ s}^{-1}$ .  
 (B) Half-life of **X** is 50 s.  
 (C) At 50 s,  $-\frac{d[\mathbf{X}]}{dt} = 13.86 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$ .  
 (D) At 100 s,  $-\frac{d[\mathbf{Y}]}{dt} = 3.46 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$ .

Q.4 Some standard electrode potentials at 298 K are given below:



To a solution containing 0.001 M of  $\text{X}^{2+}$  and 0.1 M of  $\text{Y}^{2+}$ , the metal rods **X** and **Y** are inserted (at 298 K) and connected by a conducting wire. This resulted in dissolution of **X**. The correct combination(s) of **X** and **Y**, respectively, is(are)

(Given: Gas constant,  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ ,  
Faraday constant,  $F = 96500 \text{ C mol}^{-1}$ )

- (A) Cd and Ni
- (B) Cd and Fe
- (C) Ni and Pb
- (D) Ni and Fe

Q.5 The pair(s) of complexes wherein both exhibit tetrahedral geometry is(are)

(Note: py = pyridine)

Given: Atomic numbers of Fe, Co, Ni and Cu are 26, 27, 28 and 29, respectively)

- (A)  $[\text{FeCl}_4]^-$  and  $[\text{Fe}(\text{CO})_4]^{2-}$
- (B)  $[\text{Co}(\text{CO})_4]^-$  and  $[\text{CoCl}_4]^{2-}$
- (C)  $[\text{Ni}(\text{CO})_4]$  and  $[\text{Ni}(\text{CN})_4]^{2-}$
- (D)  $[\text{Cu}(\text{py})_4]^+$  and  $[\text{Cu}(\text{CN})_4]^{3-}$

Q.6 The correct statement(s) related to oxoacids of phosphorous is(are)

- (A) Upon heating,  $\text{H}_3\text{PO}_3$  undergoes disproportionation reaction to produce  $\text{H}_3\text{PO}_4$  and  $\text{PH}_3$ .
- (B) While  $\text{H}_3\text{PO}_3$  can act as reducing agent,  $\text{H}_3\text{PO}_4$  cannot.
- (C)  $\text{H}_3\text{PO}_3$  is a monobasic acid.
- (D) The H atom of P–H bond in  $\text{H}_3\text{PO}_3$  is not ionizable in water.

**SECTION 2**

- This section contains **THREE (03)** question stems.
- There are **TWO (02)** questions corresponding to each question stem.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +2 If ONLY the correct numerical value is entered at the designated place;  
*Zero Marks* : 0 In all other cases.

**Question Stem for Question Nos. 7 and 8****Question Stem**

At 298 K, the limiting molar conductivity of a weak monobasic acid is  $4 \times 10^2 \text{ S cm}^2 \text{ mol}^{-1}$ . At 298 K, for an aqueous solution of the acid the degree of dissociation is  $\alpha$  and the molar conductivity is  $y \times 10^2 \text{ S cm}^2 \text{ mol}^{-1}$ . At 298 K, upon 20 times dilution with water, the molar conductivity of the solution becomes  $3y \times 10^2 \text{ S cm}^2 \text{ mol}^{-1}$ .

Q.7 The value of  $\alpha$  is \_\_\_\_.

Q.8 The value of  $y$  is \_\_\_\_.

**Question Stem for Question Nos. 9 and 10****Question Stem**

Reaction of  $x$  g of Sn with HCl quantitatively produced a salt. Entire amount of the salt reacted with  $y$  g of nitrobenzene in the presence of required amount of HCl to produce 1.29 g of an organic salt (quantitatively).

(Use Molar masses (in  $\text{g mol}^{-1}$ ) of H, C, N, O, Cl and Sn as 1, 12, 14, 16, 35 and 119, respectively).

Q.9 The value of  $x$  is \_\_\_\_.

Q.10 The value of  $y$  is \_\_\_\_.

### Question Stem for Question Nos. 11 and 12

#### Question Stem

A sample (5.6 g) containing iron is completely dissolved in cold dilute HCl to prepare a 250 mL of solution. Titration of 25.0 mL of this solution requires 12.5 mL of 0.03 M  $\text{KMnO}_4$  solution to reach the end point. Number of moles of  $\text{Fe}^{2+}$  present in 250 mL solution is  $x \times 10^{-2}$  (consider complete dissolution of  $\text{FeCl}_2$ ). The amount of iron present in the sample is  $y\%$  by weight.

(Assume:  $\text{KMnO}_4$  reacts only with  $\text{Fe}^{2+}$  in the solution

Use: Molar mass of iron as  $56 \text{ g mol}^{-1}$ )

Q.11 The value of  $x$  is \_\_\_\_.





Q.12 The value of  $y$  is \_\_\_\_.

## SECTION 3

- This section contains **TWO (02) paragraphs**. Based on each paragraph, there are **TWO (02)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +3 If ONLY the correct option is chosen;  
*Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered);  
*Negative Marks* : -1 In all other cases.

## Paragraph

The amount of energy required to break a bond is same as the amount of energy released when the same bond is formed. In gaseous state, the energy required for *homolytic cleavage* of a bond is called Bond Dissociation Energy (BDE) or Bond Strength. BDE is affected by *s*-character of the bond and the stability of the radicals formed. Shorter bonds are typically stronger bonds. BDEs for some bonds are given below:




Q.13 Correct match of the **C–H** bonds (shown in bold) in Column **J** with their BDE in Column **K** is

Column <b>J</b> Molecule	Column <b>K</b> BDE (kcal mol <sup>-1</sup> )
(P) <b>H</b> –CH(CH <sub>3</sub> ) <sub>2</sub>	(i) 132
(Q) <b>H</b> –CH <sub>2</sub> Ph	(ii) 110
(R) <b>H</b> –CH=CH <sub>2</sub>	(iii) 95
(S) <b>H</b> –C≡CH	(iv) 88

- (A) P – iii, Q – iv, R – ii, S – i                      (B) P – i, Q – ii, R – iii, S – iv  
 (C) P – iii, Q – ii, R – i, S – iv                      (D) P – ii, Q – i, R – iv, S – iii

Q.14 For the following reaction



the correct statement is

- (A) Initiation step is exothermic with  $\Delta H^\circ = -58 \text{ kcal mol}^{-1}$ .  
 (B) Propagation step involving  $\bullet\text{CH}_3$  formation is exothermic with  $\Delta H^\circ = -2 \text{ kcal mol}^{-1}$ .  
 (C) Propagation step involving  $\text{CH}_3\text{Cl}$  formation is endothermic with  $\Delta H^\circ = +27 \text{ kcal mol}^{-1}$ .  
 (D) The reaction is exothermic with  $\Delta H^\circ = -25 \text{ kcal mol}^{-1}$ .

### Paragraph



The reaction of  $\text{K}_3[\text{Fe}(\text{CN})_6]$  with freshly prepared  $\text{FeSO}_4$  solution produces a dark blue precipitate called Turnbull's blue. Reaction of  $\text{K}_4[\text{Fe}(\text{CN})_6]$  with the  $\text{FeSO}_4$  solution in complete absence of air produces a white precipitate **X**, which turns blue in air. Mixing the  $\text{FeSO}_4$  solution with  $\text{NaNO}_3$ , followed by a slow addition of concentrated  $\text{H}_2\text{SO}_4$  through the side of the test tube produces a brown ring.



Q.15 Precipitate X is




Q.16 Among the following, the brown ring is due to the formation of



#### SECTION 4

- This section contains **THREE (03)** questions.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +4 If **ONLY** the correct integer is entered;  
*Zero Marks* : 0 In all other cases.

- Q.17 One mole of an ideal gas at 900 K, undergoes two reversible processes, **I** followed by **II**, as shown below. If the work done by the gas in the two processes are same, the value of  $\ln \frac{V_3}{V_2}$  is \_\_\_\_.



( $U$ : internal energy,  $S$ : entropy,  $p$ : pressure,  $V$ : volume,  $R$ : gas constant)

(Given: molar heat capacity at constant volume,  $C_{V,m}$  of the gas is  $\frac{5}{2}R$ )

- Q.18 Consider a helium (He) atom that absorbs a photon of wavelength 330 nm. The change in the velocity (in  $\text{cm s}^{-1}$ ) of He atom after the photon absorption is \_\_\_\_.

(Assume: Momentum is conserved when photon is absorbed.)

Use: Planck constant =  $6.6 \times 10^{-34}$  J s, Avogadro number =  $6 \times 10^{23}$   $\text{mol}^{-1}$ , Molar mass of He = 4 g  $\text{mol}^{-1}$ )

- Q.19 Ozonolysis of  $\text{ClO}_2$  produces an oxide of chlorine. The average oxidation state of chlorine in this oxide is \_\_\_\_.

**END OF THE QUESTION PAPER**